

Design and Synthesis of Peptide-Based Nanostructures for Sensing and Therapeutic Applications

A bottom-up approach to molecular nanoscale
components

Normand Voyer

Département de chimie

Centre de recherche sur la fonction, la structure et l'ingénierie des protéines

AAAS-SWARM Meeting

April 12th, 2008




Polypeptides


Nature's choice for the construction of functional nanoscale devices



Carbonic Anhydrase



Porin




Photosynthetic Center

- Stability
- Wide variety of functional groups

Polypeptide scaffold orients functional groups in the required 3D relationship...

Different Structures: Same Function!




Chymotrypsine



Subtilisine

Modular Construction




Huge number of possible combinations to create unique nanostructures

Advantages of Polypeptides

- Choice of numerous natural and unnatural amino acids
- Well developed synthetic, purification, and characterization methods
- Ability to predict the solution 3-D structure from the primary sequence
- Allow molecular engineering

Development of Peptide-Based Molecular Devices



Development of Artificial Ion Channels

- Components in molecular sensors
- Novel therapeutic agents
- Gain fundamental insights on biological ion transport processes

Channel Proteins

- Complex membrane proteins
(MW>100,000; multiple sub-units)
- Difficult to isolate the active form
- Responsible for ion transport across cell membranes
- Extremely efficient: $>10^7$ ions/sec!
- Involved in numerous diseases


Problems with Natural Channel Proteins

- Large and complex membrane proteins
- Difficult to isolate and sensitive to denaturation
- Available in small (impure) quantities
- Very expensive
- Molecular engineering tedious

Design and Synthesis of a Family of Artificial Ion Channels


- By a simple synthetic strategy
- With possible "post-synthesis" modifications
- With different properties (ion selectivity)
- With simple monomeric active forms

KcsA: A Channel Protein Model





R. Mackinnon et al, Nature, 2001

A Minimalist Approach



KcsA



**Helical peptide
nanostructure**



**Crown ether
array**

The Prototype

N-BOC-L-CE-L₃-CE-L₂-CE-L₃-CE-L₂-CE-L₃-CE-L-OMe

- 21 Residue peptide
- Composed of L-Leucines and 6 (21-crown-7)-L-Phenylalanine (CE)
- Predicted to adopt an α -helix conformation under which the crown side chains are all aligned

Channel Design



20


16

13

9

6


2




3-4 nm

Jean-Christophe Meillon,
Normand Voyer, *Angew.
Int. Ed. Engl.*, 1997, 36, 967.

General Synthetic Strategy




Protected Segment Synthesis



Oxime Resin
 $\text{R} = \text{---C}_6\text{H}_4\text{---NO}_2$

1) TFA
2) DIEA




Segment Couplings




Activated

Heptapeptide

3) Coupling
many methods...
DIC/HOBt DMF



Final Cleavage




Desired 21-mer fully protected!

Characterization

- ^1H ^{13}C NMR Spectroscopy
- FAB, ES, MALDI Mass Spectroscopy
- Molecular Weight Range=3500-5000 g/mol
- 20 Steps
- Overall Yields=>10%
- Numerous derivatives prepared
- Available at gram scale!

Nanopore Technology









Diameters in nm


End Groups

N-terminal groups


C-terminal groups





Ion Transport Ability in Vesicles



Vesicle Studies



Membrane Conductivity



Conductivity: 2.5 pA; average open time: 1 sec (DiPhy)

Biophysical Studies

- CD spectropolarimetry (solvents, concentration, vesicles, salts, etc)
- FT~IR (solvents and vesicles)
- Polarized ATR (different peptide/lipid ration, different lipids)






Strongly α -helical, monomeric in hydrophobic solvents


Proposed Mechanism

Channel
(aggregated form in water)

+




inactive form $\Theta=90^\circ$



active form $\Theta=30^\circ$

What can we do with these compounds?



Nanoscale Therapeutics


Use synthetic channels as membrane « hole punching tools »




Wide variety of target cells




No possibility to develop resistance





Cytotoxicity Towards P388 Cancer Cells



New Approach Towards Sensing

- Develop a simple and general assay for a wide range of biologically relevant analytes

 Ultimately to the single molecule level



New Approach Towards Sensing

- Develop a simple and general assay for a wide range of biologically relevant analytes


 Ultimately to the single molecule level

- Exploit the ion transport properties of peptide nanostructures
- Exploit first the versatility of fluorescence spectroscopy

Working Hypothesis




Avidin~Biotin System



Avidin : MW = ~67 kDa

Monomer contains 128 amino acid residues




Biotin : MW = 244.31




$K_a = 10^{15} \text{ M}^{-1}$



Biotinylated Nanostructures





1




3




2




4



Fluorescence Studies



Control Experiments



Conclusions

- Novel and general approach to artificial ion channels
- Compounds adopt the predicted α -helix conformation in TFE and in membranes
- Exist as monomers in low polarity environment

Conclusions (Part 2)

- Function as true artificial ion channels in vesicles and in bilayer membranes
- Ions travel by the channel formed by the crown ethers alignment
- Cytotoxic to cancer cells

Conclusions

- Developed a fluorescent assay using peptide nanostructures as « molecular transducers »
- Simple and applicable to a wide range of clinically important label-free analytes

Acknowledgments

Graduate Students

M.E. Provencher

J.-P. Blanchette

M. Arseneault

J. Richer

J. Martin-Boutin

M. Ouellet

D. Hivon

M. Dumont

P.-L. Boudreault

S. Ouellet

J.-C. Meillon

E. Biron

M.Tremblay*

F. Otis*

M. Beaumont

M. Giguère

C. Barberis

J. Roby

M. Robitaille

J. Lamothe

B. Guérin

F. Maltais

A. Pichette

Collaborators

M. Auger (Laval)

B. Smith (UND)

B. Pucci (Avignon)

G. Dewynter (Montpellier)

R. Gaudreault (Laval)

Postdocs

J. Joutel

R. Barratin

Financial \$\$\$upport

- CRSNG/NSERC of Canada
- NanoQuébec
- FQRNT Gouvernement du Québec
- FRSQ Gouvernement du Québec
- Université Laval
- Ministères de l'enseignement supérieur
France/Québec



The Québec Advantage*

8 Months without air conditioning needs



2008 Québec City 400th

An



Thank you!




Comparison

21-Crown-7 channel 18-Crown-6 channel


Functional	Yes	Yes
Conductance	2.5 and 5 pA	3.5 pA
Lifetime	>1 sec	100-200 msec

Circular Dichroism Studies



Different solvents...



Different lengths...




CD in Vesicles




[] = 10⁻⁵ - 10⁻³ M in TFE

EYL Vesicles lipid/peptide=80/1

FTIR Spectroscopy



ATR Studies



Using polarized ATR spectroscopy, the angle θ between a helix and the bilayer normal can be found...

$$f(\theta) = \frac{R - 2}{R + 1.45} \frac{2}{3 \cos^2 \theta - 1}$$


$$f(\theta) = \frac{3 \cos^2 \theta - 1}{2}$$

An arrow points from the θ term in the denominator of the second equation to the θ term in the first equation.

ATR Results

Samples	Spectral bands			
	$V_{\text{CH}_2 \text{ sym}}$		$V_{\text{amide 1}}$	
	R	θ	R	θ
DMPC	1.09±0.01	25.5±0.1		
BOC-21 mer (21-C-7)-Me 1	1.14±0.01	28.3±0.2	1.84±0.02	58.7±0.2
BOC-21 mer (18-C-6)-OMe 2	1.14±0.01	28.3±0.2	1.76±0.20	59.4±1.2
BOC-21 mer (13-C-4)-OMe 3	1.18±0.02	29.9±1.0	2.01±0.05	54.6±0.9
BOC-21 mer (13-C-4)-OH 4	1.17±0.02	29.5±1.0	1.86±0.02	57.5±0.3
H-21 mer (13-C-4)-OH 5	1.29±0.01	34.3±0.3	1.84±0.01	57.6±0.1

Towards Microarray Nanosensors



Gold patterned microarray

Simultaneous detection of analytes
at single molecule level